

**Composition :** Ciprozid-XR 1000 : Each extended release tablet contains Ciprofloxacin 1000mg as Ciprofloxacin Hydrochloride USP.

**Pharmacology :** Ciprofloxacin is a synthetic 4-quinolone derivative with bactericidal activities against a wide range of gram-negative and gram-positive organisms. The absolute bioavailability of Ciprofloxacin when given as an oral tablet is approximately 70% with no substantial loss by first pass metabolism. Maximum serum concentrations are attained 1 to 2 hours after oral dosing.

**Indications :** Ciprozid-XR 1000 is indicated only for the treatment of urinary tract infections, including, Uncomplicated urinary tract infections (Acute cystitis) caused by Escherichia coli, Proteus mirabilis, Enterococcus faecalis or Staphylococcus saprophyticus; Complicated urinary tract infections caused by Escherichia coli, klebsiella pneumoniae, Enterococcus faecalis, Proteus mirabilis or Pseudomonas aeruginosa; Acute uncomplicated Pyelonephritis caused by Escherichia coli.

**Dosage and Administration :** Ciprozid-XR 1000 should be administered orally once daily in the following dosage guideline table.

#### Dosage Guidelines

Indication	Unit Dose	Usual Durations
Uncomplicated Urinary Tract Infection (Acute Cystitis)	500mg	3 Days
Complicated Urinary Tract Infection	1000mg	7-14 Days
Acute Uncomplicated Pyelonephritis	1000mg	7-14 Days

Or, as directed by the registered physician.

**Contraindications :** This medicine is contraindicated to patients with hyper sensitivity to any component of this product

**Ciprozid-XR 1000  
Tablet**



**DRUG  
INTERNATIONAL  
LTD.**

or, any member of quinolone class of antimicrobial agents.

**Precautions :** It should be used with caution in epileptics and patients with a history of CNS disorder.

**Side effects :** Adverse effects include nausea, diarrhea and other gastrointestinal disturbances, headache, dyspepsia, dizziness, skin rashes, vomiting etc.

**Use in pregnancy and lactation :** There are no adequate and well controlled studies in pregnant women. Ciprofloxacin is excreted in human milk so, it should be used if clearly needed.

**Use in Child :** There is no data available.

**Drug Interactions :** Ciprozid-XR 1000 should be avoided with antacids or sucralfate or other products containing iron and multivitamin preparation with zinc.

**Overdose :** There is no data available.

**Storage :** Store below 30°C in a dry place.

**Packing :** Ciprozid-XR 1000 : Each box contains 14's tablets in blister pack.