

Composition : Each ml sterile Eye & Ear Drops contains Ciprofloxacin USP 3 mg as Ciprofloxacin hydrochloride.

Indications : Eye : Ciprofloxacin is indicated for the treatment of corneal ulcers and superficial infections of the eye and adnexa caused by susceptible strains of bacteria. **Ear :** Otitis Externa.

Dosage and administration : Eye : Adults : Superficial ocular infection: The usual dose is one or two drops in the affected eyes four times a day. In severe infections, the dosage for the first two days may be one or two drops every two hours during waking hours.

Bacterial corneal ulcer : Two drops into the affected eye every 15 minutes for the six hours and then two drops into affected eye every 30 minutes for the remainder of the first day. On the second day, instill two drops in the affected eye hourly. On the third to the fourteenth day, place two drops in the affected eye every four hours. Treatment may be continued after 14 days if corneal reepithelialization has not occurred. **Children :** The safety & efficacy of Ciprofloxacin in children under the age of 11 years has not been established. **Ear :** One or two drops is instilled into the infected ear every four hours. Or, as directed by the registered physician.

Contraindication : A history of hypersensitivity to Ciprofloxacin or any other component of the medication is a contraindication to its use.

Ciprozid
Eye & Ear Drops



DRUG
INTERNATIONAL
LTD.

Precaution : As with other antibacterial preparations, prolonged use of Ciprozid ophthalmic solution may result in overgrowth of nonsusceptible organisms, including fungi. If superinfection occurs, appropriate therapy should be initiated. Ciprofloxacin should be discontinued at the first appearance of a skin rash or any other sign of hypersensitivity.

Side effects : The most frequently reported drug related adverse reaction was local burning or discomfort. Other reactions included lid margin crusting, crystals/scales, foreign body sensation, itching, conjunctival hyperemia and a bad taste following instillation.

Use in Pregnancy and Lactation : There is no adequate and well-controlled study in pregnant women. Ciprofloxacin ophthalmic solution should be used during pregnancy only if the potential risk to the fetus. Caution should be exercised when Ciprofloxacin ophthalmic solution is administered to a nursing mother.

Drug Interactions : Specific drug interaction studies have not been conducted with ophthalmic Ciprofloxacin.

Packing : 10ml in plastic dropper bottle.