

Composition : Zolmitriptan USP 2.5mg Tablet.

Indications : Zolmitriptan is indicated for the acute treatment of migraine with or without aura in adult.

Dosage and administration : The recommended dose of Zolmitriptan to treat a migraine attack is 2.5mg. If symptoms persist or return within 24 hours, a second dose has been shown to be effective. If a second dose is required, it should not be taken within 2 hours of the initial dose. If a patient does not achieve satisfactory relief with 2.5mg doses, subsequent attacks can be treated with 5mg doses of Zolmitriptan. In the event of recurrent attacks, it is recommended that the total intake of Zolmitriptan in a 24 hour period should not exceed 15mg. Or, as directed by the registered physician.

Contraindications : Zolmitriptan is contraindicated in patients with : ● Known hypersensitivity to any component of the product. ● Uncontrolled hypertension. ● Ischaemic heart disease. ● Coronary vasospasm / Prinzmetal's angina.

Side effects : Zolmitriptan is well tolerated. Adverse reactions are typically mild/moderate, transient, not serious and resolve spontaneously without additional treatment. Possible adverse reactions tend to occur within 4 hours of dosing and are no more frequent following repeated dosing. The following adverse

Miotrol
Tablet



**DRUG
INTERNATIONAL
LTD.**

somnolence; warm sensation; asthenia; dry mouth.

Precautions : Care should be taken to exclude other potentially serious neurological conditions. zolmitriptan should not be given to patients with symptomatic Wolf-Parkinson-White syndrome or arrhythmias.

Pregnancy and lactation : Pregnancy Category C by FDA.

Drug interactions : There is no evidence that concomitant use of migraine prophylactic medications has any effect on the efficacy or unwanted effects of Zolmitriptan.

Packing : Miotrol : 5 x 10's tablets in blister pack.