

Composition : **Canarol-100 Tablet** : Each film coated tablet contains Canagliflozin 100 mg as Canagliflozin Hemihydrate INN.

Indications : Canagliflozin is indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus.

Dosage and administration : 100 mg tablet is taken orally before the first meal of the day. Or, as directed by the registered physician.

Contraindications : It is contraindicated in patients having serious hypersensitivity reaction to the Canagliflozin or end stage renal disease or patients on dialysis.

Precautions : Canagliflozin is not for treatment of type 1 diabetes or diabetic ketoacidosis. It should not be used to patient on dialysis or having severe kidney disease.

Side effects : The most common side effects are female genital mycotic infections, urinary tract infections and increased urination.

Pregnancy and lactation : There are no adequate and well-controlled studies of Canagliflozin in pregnant women. It is unknown whether Canagliflozin is secreted in human breast milk. Therefore, it should only be used to pregnant

Canarol
Tablet



women & nursing mothers only if the potential benefit justifies the potential risk to the fetus.

Drug Interactions : Co-administration of Canagliflozin with non-selective inducer of several UGT enzymes (e.g., Rifampin, Phenytoin, Phenobarbital, Ritonavir) decrease its efficacy. There was an increase in the AUC and Cmax of digoxin when co-administered with Canagliflozin tablet.

Overdose : There were no reports of overdose during the clinical development program of canagliflozin.

Storage : It should be stored at 20-25°C away from heat, moisture and light. Keep out of reach of children.

Packing : Canarol-100 Tablet : Each box contains 1 x 10's tablets in blister pack.