CARDIZEM-SR-

Diltiazem Hydrochloride USP

Composition: Cardizem-SR 90: Each sustained release tablet contains 90mg of Diltiazem Hydrochloride USP.

Cardizem-SR 120 : Each sustaine d release tablet contains 120mg of Diltiazem Hydrochloride USP.

Description : Cardizem-SR (Diltiazem HCI) is a calcium antagonist. It selectively reduces calcium entry through voltage-dependent calcium channels into vascular smooth muscle cells and myocardial cells. This lowers the concentration of intracellular calcium. This action of diltiaze m results in dilation of coronary arteries causing an increase in myocardia I oxygen supply. It reduces cardia c work by moderating the heart rate and reducing systemic vascular resistanc e thus reducing oxygen demand.

Indications : 1. Mild to moderate cases of hypertension. 2. Angina pectoris

Dosage and administration : Dosage must be adjusted accordin g to each patient's needs. Mild to moderate cases of hypertension: Initially 90mg or 120mg twice daily (elderly once daily); upto 360mg daily may be required (elderly upto 240mg daily) Relief of Anginal Pain: Initially 90mg or 120mg twice daily in divided doses may be required (elderly upto 240mg daily). Or, as directed by a registered physician.

Contraindications: Cardizem-SR is contraindicated in:

- 1. Patients with sick sinus syndrome except in the presence of a functioning ventricular pace maker.
- 2. Patients with 2nd or 3rd degree AV block or sinoatrial block except in the presence of a functioning ventricular pace maker.
- 3. Patients who have demonstrated hypersensitivity to the drug.
- 4. Pregnant women or women suspected of being pregnant.
- 5. Patients with acute myocardial infarction and pulmonary congestion documented by X-ray.

6. Patients with hypotension.

Precautions : 1. Intake of Cardizem -SR to be reduced gradually, not to be withdrawn suddenly or without doctor 's advice.

- 2. Patients with severe bradycardia or 1st degre e AV block.
- 3. In case of impaired renal & hepatic functions, in case of dermatologic reactions diltiazem should be discontinued.

Side effects: Bradycardia, sino-atrial block, atrioventricular block, hypertension malaise, headache, hot flushes. GIT disturbances, oedema, hepatitis and depression

Drug interactions : Cardizem-SR should be carefully administered in case of concomitant use with the following drugs:

1. Anti hypertensive agent. 2. Beta blocking agent. 3. Carbamazepine. 4. Digoxin preparations. 5. Cimetidine, 6. Anesthetics.

Storage: Store below 30 °C in a dry place.

Packing : Cardizem-SR 90 : Each box contains 5 x 10 's tablet in blister pack.

Cardizem-SR 120: Each box contains 5 x 10 's tablet in blister pack

DRUG INTERNATIONAL LTD.

05/2023/Nil