

Composition: Each Tablet Contains Deucravacitinib INN 6mg.

Pharmacology: The absolute oral bioavailability of Deucravacitinib was 99% and ranged from 2 to 3 hours in healthy patients. The volume of distribution at steady state is 140 L. The terminal half-life is 10 hours. The renal clearance ranged from 27 to 54 mL/Minute. Deucravacitinib is metabolized by cytochrome P-450 (CYP) 1A2. Approximately 13% and 26% of the Deucravacitinib dose excreted by urine and feces.

Indication: It is indicated for the treatment of adults with moderate-to-severe plaque psoriasis who are candidates for systemic therapy or phototherapy.

Dosage & Administration: The recommended dosage is 6mg taken orally once daily, with or without food. Or, as directed by a registered physician.

Contraindications: It is contraindicated in patients with known hypersensitivity to Deucravacitinib or any other components of this product.

Precautions: Infections: It may increase the risk of infection. Avoid use in patients with active or serious infection. **Tuberculosis:** Evaluate for TB prior to initiating treatment with this. **Malignancy:** Malignancies

Decanib 6 Tablet



DRUG
INTERNATIONAL
LTD.

including lymphomas were observed in clinical trials.

Side effects: Most common side effects are upper respiratory infections, blood creatine phosphokinase increased, herpes simplex, mouth ulcers, folliculitis and acne.

Use in Pregnancy and Lactation:

Pregnancy: There is no data available. **Lactation:** There are no data on the presence of Deucravacitinib in human milk, the effects on the breastfed infants, or the effects on milk production.

Use in Child: The safety and effectiveness of Deucravacitinib in pediatric patients have not been established.

Drug-Interactions: Drug interaction of deucravacitinib was observed with following drugs: Cyclosporine, Ritonavir, Famotidine or Rabeprazole.

Overdose: There is no data available.

Storage: Store below 30°C away from sunlight. Keep out of reach of children.

Packaging: Each box contains 30's tablet in blister pack.

