

Composition : Each ml sterile ophthalmic suspension contains Fluorometholone USP 1 mg.

Indications : It is indicated for the treatment of corticosteroid-responsive inflammation of the palpebral and bulbar conjunctiva, cornea and anterior segment of the globe.

Dosage and administration : One drop is instilled into the conjunctival sac two to four times daily. During the initial 24 to 48 hours, the dosing frequency may be increased to one application every four hours. Care should be taken not to discontinue therapy prematurely. Or, as directed by the registered physician.

Side effects : Adverse reactions include elevation of intraocular pressure with possible development of glaucoma and infrequent optic nerve damage, posterior subcapsular cataract formation, delayed wound healing and perforation of the globe.

Contraindications : It is contraindicated in individuals with known or suspected hypersensitivity to any of the ingredients of this preparation and to other corticosteroids.

Use in pregnancy and lactation : There are no adequate and well-controlled studies of fluorometholone in pregnant women. Fluorometholone should be used during pregnancy only if the potential benefit Justifies the potential risk to the fetus. It is not known whether topical ophthalmic administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in human milk.

F-Lon
Ophthalmic Suspension



**DRUG
INTERNATIONAL
LTD.**

Because of the potential for serious adverse reactions in nursing infants from fluorometholone, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Precautions : As fungal infections of the cornea are particularly prone to develop coincidentally with long-term local corticosteroid applications, fungal invasion should be suspected in any persistent corneal ulceration where a corticosteroid has been used or is in use. If this product is used for 10 days or longer, intraocular pressure should be monitored.

Drug interactions : No information provided.

Overdose : No data are available in regard to overdosage in humans.

Storage : Store below 25oC away from sunlight.

Packing : Each box contains 5ml in a plastic dropper bottle.